THE UNIVERSITY OF NEW MEXICO
HEALTH SCIENCES CENTER

Insulin Order Set Administration Record

Allergies:

Patient Label

Nursing Orders if NPO:

1. Check blood glucose every 4 hours at: 2400 – 0400 – 0800 - 1200 – 1600 – 2000.
2. Give "Correction Dose" insulin based on Rapid-Acting Insulin table
3. Administer insulin after each CBG per table, do not recheck CBG for 4 hours
4. Do not hold glargine for NPO status
5. Do not mix glargine in same syringe with other insulins

Nursing Orders if Eating:

1. Check blood glucose just before the patient eats meals and at 2200
2. Calculate and give "Nutritional + Correction Dose" insulin based on Rapid-Acting Insulin table
3. Give only correction doses at bedtime and if the patient eats less than 50% of their meal
4. Administer the insulin within 30 minutes of first bite of the meal
5. Do not mix glargine in same syringe with other insulins

Blood Glucose mg/dL	Rapid Acting Insulin Table Type of Insulin: _____	
≤ 70	Implement Hypoglycemia orders 1. If patient can take po: administer 15 grams of carbohydrate (6 oz of fruit juice or 15 grams of glucose gel) 2. If patient unable to take po: administer 12.5 grams of dextrose 50% IV (25 mL) or 1 mg of glucagon IM 3. Check blood glucose every 15 minutes and repeat above treatment until blood glucose is ≥ 100 mg/dL 4. Notify MD if blood glucose ≤ 70 mg/dL for two readings in 24 hour period	
	Correction Dose	**Nutritional Dose**
71-149	+ _____ units	___ units with Breakfast ___ units with Lunch ___ units with Dinner
150-199	+ _____ units	
200-249	+ _____ units	
250-299	+ _____ units	
300-349	+ _____ units	
350-399	+ _____ units	
≥ 400	Treat with +_____ units and inform MD	

Date / /		Date / /		Date / /		Date / /		Date / /	
Time	CBG								
Initial x2	Tx								

Date	Signatures	Date	Signatures	Date	Signatures

Diagnosis:
Admission Date:
Today's Date:
Physician:
Sex:
Height:
Weight:

Please contact krogers@salud.unm.edu with any concerns, questions, or suggestions on this order set.

Revised 9/2007